

Comparison of blood pressure reduction between treatments of steeping Red Ginger Rhizome and steeping Binahong Leaves

by Djelang Fickri

Submission date: 20-Aug-2021 01:11PM (UTC+0700)

Submission ID: 1633541223

File name: 16304-Article_Text-47741-1-2-20210418.docx (80.21K)

Word count: 4267

Character count: 23343



Research Article

Comparison of blood pressure reduction between treatments of steeping Red Ginger Rhizome (*Zingiber officinale* R.) and steeping Binahong Leaves (*Anredera cordifolia* (Ten.) Steenis) in Healthy People with hypertension risk

Djelang Z. Fickri^{(1)*}, Arista W. Ningsih⁽²⁾, Martina K. Rohmah⁽¹⁾, Khurin I. Wahyuni⁽¹⁾, Novita E. Winarni⁽¹⁾, Pahri⁽¹⁾

(1) Clinical Pharmacy department, High School of Health Sciences, Anwar Medika Hospital, Sidoarjo Indonesia

(2) Biological Pharmacy department, High School of Health Sciences, Anwar Medika Hospital, Sidoarjo Indonesia

* Corresponding author's email : fickri_farma@yahoo.co.id

ARTICLE INFO

ABSTRACT

Article History

Submitted MMDD,YY
 Revised MMDD, YY
 Accepted, MMDD, YY
 Published, MMDD, YY

Keywords

Anredera cordifolia (Ten.) Steenis
 Binahong
Zingiber officinale R.
 Red Ginger
 Hypertension

doi:

10.22219/farmasains.XX.YYYY

Background: The potency of red ginger rhizome and binahong leaves has been known have antihypertensive effect and can be used in herbal medicine. This study was to compare the effect of steeping red ginger rhizome and steeping binahong leaves on blood pressure reduction in healthy people with hypertension risk. **Methods:** The research design is Comparative Quasy Experiment to measure pre-test and post-test blood pressure on treatment of 3 g / 200 ml steeping the red ginger rhizome and 3gr / 200 ml steeping binahong leaves. The number of respondents was 64 according to the inclusion criteria and had obtained the ethical clearance. The data were processed statistically. **Results:** The experiment showed that the S1 group had a significant difference in systolic blood pressure reduction compared with the control treatments ($P < 0.05$), but there was no significant difference in diastolic blood pressure reduction compared with controls ($P = 0.097 > 0.05$). Meanwhile, in the S2 group, there was a significant difference in systolic blood pressure reduction compared with the control treatments ($P < 0.05$), but there was no significant difference in diastolic blood pressure reduction compared with controls ($P = 0.512 > 0.05$). Between the treatment of steeping red ginger rhizome (S1) and steeping binahong leaves (S2) there was no significant difference in systolic blood pressure reduction, which means that both red ginger and binahong seed had the same effect on systolic blood pressure ($p = 0.78 > 0.05$). **Conclusions:** Treatment steeping red ginger rhizome (*Zingiber officinale* R.) and steeping binahong leaves (*Anredera cordifolia* (Ten.) Steenis) has a

similar effect on systolic blood pressure reduction.

1. Introduction (Calibry, body 12 pt, single space)

Hypertension is defined as persistent blood pressure where the systolic and diastolic blood pressure is above 140/90 mmHg and above 160/90 mmHg in the elderly [1]. According to Yonata and Pratama, if hypertension can be properly controlled, it can reduce the risk of death, cardiovascular disease, stroke, and heart failure [2]. According to WHO data, approximately 26.4% of the world's population has hypertension, and it is likely that it will continue to increase to 29.2% in 2025 if the condition of the people's lifestyle is not controlled [3]. Hypertension is a progressive disease and has become a major global health problem. Of the 972 million people with hypertension, 333 million are at in developed countries and 639 million are in developing countries including Indonesia [2]. In general, the management of hypertension is pharmacological and non-pharmacological. Pharmacological therapy includes chemical and herbal medicines. Pharmacological treatment with chemical drugs have adverse effects such as the use of thiazide-type diuretics in hypertensive patients accompanied by hyperuricemia can lead to joint inflammation. Hypertensive patients with a history of asthma should avoid β -blocker as it can aggravate asthmatic conditions [4]. So, it is necessary to develop the potential treatment from natural ingredients because it has minimal side effects.

Alternative hypertension treatment from natural ingredients, for example, is red ginger and binahong [5, 6]. The potency of the red ginger rhizome and binahong leaves has been known to have an antihypertensive effect and can be used in herbal medicine. In previous studies, the antihypertensive effect of 70% ethanol extract from red ginger was tested, which had a significant effect on systolic blood pressure reduction compared with negative controls, and the highest systolic blood pressure reduction was 27.35% in rats [7]. The main bioactive compounds in red ginger are 4-diarylheptanoid, shogaol, gingerol, and gingeron, all of which have higher antioxidant activity than vitamin E [8]. In another in vivo test, red ginger has antihypertensive activity in salt-induced mice for 5 weeks [9]. Gingerol is known to reduce blood pressure systemically in mice at dose 10–100 $\mu\text{g}/\text{kg}$ and cause bradycardia when given intravenously [10, 11]. Other studies have shown that Gingerol in ginger have anticoagulant effect, which prevents blood clots, thereby preventing blockage of blood vessels, which is the main cause of strokes and heart attacks [12]. (6)-shogaol and (6)-gingerol, (10)-gingerol have antioxidant effects [13]. Antioxidants can reduce free radicals such as superoxide anion, thromboxane A₂, endothelin, and endoperoxide which can cause hypertension. Superoxide anions can reduce nitric oxide. While tromboxane A₂, endothelins, and endoperoxides a trigger factor endothelial vasoconstriction. Antioxidants are able to increase the formation of nitric oxide (NO) [14]. Nitric oxide (NO) causes vasodilation events [15]. At the time of vasodilation, Total Peripheral Resistance (TPR) decreases, resulting in a lowering of blood pressure. Flavonoids have an inhibitory effect that prevents the activity of angiotensin-converting enzyme (ACE) [16]. Flavoniod prevents ACE activity, causing vasodilation, then decrease in cardiac output and ultimately decrease in blood pressure [17].

Binahong leaves are empirically often used routinely as many as 9 pieces every day as a blood pressure lowering. The results of the research by Astuti et al., showed that binahong leaves contain flavonoids, alkaloids, saponins, steroids, triterpenoids, and polyphenols [18]. The flavonoid content in fresh samples of binahong leaves was 11.23 mg / kg and in dry samples was 7.81 mg / kg [19]. In research Hanafi et al, leaves binahong have averaged 1.08% of saponins, 1.58% of tannins and 0.39% of flavonoids [20]. The flavonoids in binahong leaves allow the plant to be used to reduce blood pressure. Flavonoids may inhibit ACE, so that will

inhibit the conversion of angiotensin I to angiotensin II and decrease the secretion of antidiuretic hormone. As a result, a lot of urine is excreted and eventually the blood pressure decreases [21]. Type of flavonoids compound in extracts binahong are flavonols. These compounds have the ability to eliminate oxidative stress, inhibit the activity of angiotensin converting enzyme (ACE), improve endothelial relaxation of blood vessels, regulate cell signaling and gene expression. Flavonols act as major antioxidants in biological systems. Flavonols show potential as ACE inhibitor in vitro and in vivo [22]. Aidasari's research states that the ethanol extract of binahong leaves is effective in reducing the blood pressure of male Wistar rats at a dose of 50 mg / kgBB ($p = 0.05$) and at a dose of 100 mg / kgBW ($p = 0.002$) [23]. According to research by Siswantari et al., giving binahong leaves boiled water can lower blood pressure in the elderly with the t-test value of 3.873 (df 14, $p = 0.002$) [24].

Both of these herbal ingredients have the active compounds is believed to be antihypertensive with almost the same mechanism of vasodilator. Necessary to study both the herbal ingredient if administered in the form of steeping water. The effect of lowering blood pressure was assessed as a direct response to the administration of steeping the red ginger rhizome and steeping binahong leaves.

2. Materials and Methods

Research design

The research design is Comparative Quasy Experiment to measure the pre-test and post-test blood pressure on treatment of 3 g / 200 ml steeping the red ginger rhizome and 3gr / 200 ml steeping binahong leaves, measuring pre-test and post test with a range of 4 hours. Both groups were observed before treatment, then observed again after treatment. Subjects were observed in blood pressure (pretest), then given a treatment in the group of Red Ginger (S1) and the group leaves Binahong (S2), then observed to measure blood pressure (posttest) after 4 hours. The independent variable was the treatment of steeping the red ginger rhizome 3 g / 200 ml and binahong steeping leaves 3 g / 200 ml and the dependent variable is blood pressure value.

Population and Sample

Population in the study were people in Desa Bungkeng, Kecamatan Tanjung Bumi, Kabupaten Bangkalan - Madura who were given steeping red ginger rhizome (S1) and people in Desa Gempol Kurung, Kecamatan Menganti, Kabupaten Gresik - East Java who were given steeping binahong leaves (S2). The number of samples was determined by the Lameshow formula and using non random purposive sampling method. Samples according to the inclusion and exclusion criteria of the study.

The inclusion criteria in this study included: over 25 years of age, obesity (BMI > 25), smokers, excessive caffeine consumption, excess salt consumption. The exclusion criteria included: having a history of allergies to steeping binahong leaves and steeping red ginger, disliking steeping red ginger and binahong steeping, taking hypertension drugs or other drugs.

Treatment Preparation

The process of steeping red ginger rhizome and steeping binahong leaves is as follows: Clean the red ginger and binahong leaves first. After washing, it is chopped small and dried by peeling it at room temperature. After that, it is crushed into powder. 3 grams of dry red ginger powder is put into a tea pouch. By the same token with binahong leaves powder. Then steeped with 200 ml hot water at 90°C until the water color change and let stand for 2 minutes. By the same token with binahong leaves powder.

The process of preparing a placebo is the same as the method above and filling it with 3 grams small chunks of tea pouch and put into the tea pouch. Then steeped with 200 ml hot water at 90°C until the water color change and let stand for 2 minutes.

Data analysis method

Statistical analysis using the SPSS 22.0 program. Data were tested for normality with a significance level of $\alpha = 0.05$. Paired T test was performed to test the differences in each group on parametric data, and Wilcoxon test on non-parametric data with a significance level of $\alpha = 0.05$. The comparison test used the Mann Whitney test with a significance level of $\alpha = 0.05$.

Ethical approval

Research involving healthy people and complied with all relevant national regulations and institutional policies with Faculty of Dental Medicine Health Research Ethical Clearance Commission (Faculty of Dental Medicine, Universitas Airlangga, Surabaya, Indonesia) was studied the reseacrh design carefully, was declare to be ethically appropriate in accordance to 7 (seven) WHO 2011 and was certified (100/HRECC.FODM/III/2020 & 196/HRECC.FODM/IV/2020) for the care and use of human subjects.

3. Results and Discussions

Result

Characteristics of the respondents are presented in the table below:

Tabel 1. Respondent characteristics

Category	S1 ¹		S2 ²	
Gender				
- Male	17	53%	12	37,5%
- Female	15	47%	20	62,5%
N	32	100%	32	100%
Age				
- 25-35 years	13	40%	15	46,9%
- 36-45 years	6	19%	14	43,7%
- 46-55 years	8	25%	3	9,4%
- 56-65 years	5	16%		
N	32	100%	32	100%
Jon				
- private employees	8	25%	12	37,5%
- entrepreneur	2	6%	5	15,6%
- housewife	7	22%	11	34,4%
- farmer	15	47%	4	12,5%
N	32	100%	32	100%
The habits can be a risk factor for hypertension.	Eating excessively salty, smoking cigarettes, drinking coffee		Eating excessively salty, spicy food, drinking coffee	

¹S1 : People in Desa Bungkeng, Kecamatan Tanjung Bumi, Kabupaten Bangkalan - Madura who were given steeping red ginger rhizome

²S2 : People in Desa Gempol Kurung, Kecamatan Menganti, Kabupaten Gresik - East Java who were given steeping binahong leaves

The number of respondents was 64 people consisting of two groups from different locations, from Desa Bungkeng (S1) and from Desa Gempol Kurung (S2), and each of these groups had 2 treatment groups with the

same individual being a negative control group and treatment group. In the negative control group, respondents were given placebo intervention, while in the treatment group the respondents were given steeping red ginger (S1) and steeping binahong leaves (S2).

Description of data will show mean value, standard deviation, maximum and minimum value. Description of data are shown in the following table:

Table 2. Pre-test and post-test blood pressure value in the control groups

Group	Variable	N	Maks	Min	Mean ± SD	P	
Control (S1)	Systolik	Pre-test	32	214	101	140 ± 27	0,620
		Post-test	32	212	102	141 ± 26	
	Diastolic	Pre-test	32	144	60	90 ± 17	0,092
		Post-test	32	142	67	92 ± 16	
Control (S2)	Systolik	Pre-test	32	157	131	144 ± 7	0,720
		Post-test	32	156	130	142 ± 8	
	Diastolic	Pre-test	32	96	85	92 ± 3	0,600
		Post-test	32	94	81	90 ± 3	

Wilcoxon signed rank test ($\alpha = 0,05$)

Table 3. Pre-test and post-test blood pressure value in the treatment groups

Group	Variable	N	Maks	Min	Mean ± SD	P	
S1	Systolik	Pre-test	32	199	99	141 ± 27	0,000
		Post-test	32	182	100	134 ± 16	
	Diastolic	Pre-test	32	148	65	91 ± 24	0,000
		Post-test	32	143	62	87 ± 17	
S2	Systolik	Pre-test	32	162	133	148 ± 8	0,000
		Post-test	32	156	123	142 ± 8	
	Diastolic	Pre-test	32	100	80	92 ± 3	0,000
		Post-test	32	100	80	90 ± 3	

Wilcoxon signed rank test ($\alpha = 0,05$)

Table 4. Blood pressure reduction value and compared with controls

Group	Variable	N	Maks	Min	Mean ± SD	P	
S1	Systolik	Control	32	5	2	3,23 ± 1,23	0,001
		Treatment	32	8	1	4,32 ± 1,08	
	Diastolic	Control	32	6	3	3,24 ± 1,43	0,097
		Treatment	32	9	3	5,43 ± 1,78	
S2	Systolik	Control	32	5	1	2,44 ± 1,24	0,000
		Treatment	32	10	2	5,25 ± 1,95	
	Diastolic	Control	32	4	0	1,44 ± 1,08	0,512
		Treatment	32	5	0	1,75 ± 1,49	

Wilcoxon signed rank test ($\alpha = 0,05$)

Table 5. Comparison of Blood Pressure Reduction Value of S1 and S2 Respondents

Variable	Mean \pm SD		P
	S1	S2	
Systolic	4,32 \pm 1,08	5,25 \pm 1,95	0,078
Diastolic	5,43 \pm 1,78	1,75 \pm 1,49	0,000

Mann whitney U test ($\alpha = 0,05$)

Discussion

All respondents did not have a history of hypertension diagnosis and looks like a healthy people even though most respondents had high blood pressure. Men are more susceptible to hypertension because of a worse lifestyle by smoking and drinking coffee which can cause increase blood pressure due to the nicotine in cigarettes and caffeine in coffee. The risk of hypertension increases with age. It is caused by changes in the structure and function of peripheral vascular system that is responsible for changes in blood pressure. These changes include the formation of atherosclerosis, loss of fibroblast elasticity and decreased smooth muscle relaxation. As a result, the aorta and arteries decrease in stretch and decrease the capacity to accommodate blood volume from the heart. then lead to increase in cardiac output and increase in peripheral resistance so that blood pressure will rise [16, 17].

pressure and stressful job demands can stimulate the release of adrenal hormones and trigger the heart to beat faster, so that the blood pressure will rise. Anxiety at work will affect the sympathetic nerve activity and vasoconstriction occurs. Vasoconstriction resulting in renal blood flow down and can cause renin release. Renin stimulates the formation of angiotensin I which is then converted into angiotensin II. This hormone causes sodium and water retention in the renal tubules so that intravascular pressure increases and consequence in increased blood pressure [16, 17]. Respondents almost never felt the symptoms of hypertension in direct interviews.

Sodium intake, stress, obesity, genetics, can affect cardiac output and peripheral resistance. Blood pressure will increase if there are abnormalities in these factors and will result in changes in the walls of the aorta and blood vessels. Furthermore, the systolic blood pressure increases with no or little diastolic changes. Increased systolic blood pressure causes the heart workload become heavier and eventually the walls of left ventricle thicken due to adaptation. Initially, the thickening of verticle wall is for adaptation, over time will increase heart workload and become pathological condition. When heart workload increases, the heart will pump blood faster and blood pressure will rise.

In this study, all male respondents had the habit of smoking and drinking coffee and almost all of the female respondents had a salty appetite or high sodium consumption which resulted in high blood pressure (Table 1). Excess salt intake can cause fluid buildup in the body because it attracts extracellular fluid, which increases blood volume and pressure. This proves that the respondent's food intake will affect blood pressure. The smoking habit will damage the endothelial lining of blood vessels due to nicotine and carbon monoxide from cigarettes, which then triggers the atherosclerosis process and blocks the supply of oxygen and nutrients to organs, especially the heart. The presence of atherosclerosis will increase cardiac output and blood pressure [4]. While at the coffee drinking habits with the largest content of caffeine will bind adenosine receptors, activate the sympathetic nervous system by increasing concentrations of catecholamines in plasma and increases the production of cortisol. It impacts on vasoconstriction and increase in total peripheral resistance which will cause an increase in blood pressure [16, 17].

The control group S1 respondents as many as 20 respondents observed a decrease in systolic pressure and 23 respondents observed a decrease in diastolic blood pressure ($p = 0.620 > 0.05$ & diastolic $p = 0.092 > 0.05$), which means that giving placebo did not provide a significant change in blood pressure (Table 2). While the treatment group of 30 respondents observed a decrease in systolic pressure and 28 respondents observed a decrease in diastolic blood pressure (systolic $p = 0.000 < 0.05$ & diastolic $p = 0.000 < 0.05$), which means that the treatment of steeping red ginger gives a significant change in blood pressure (Table 3).

Red ginger has benefits in the cardiovascular system. Red Ginger has antioxidant effects that can reduce free radicals and can reduce blood pressure through calcium channel blockade [13]. Red ginger can also cause a decrease in blood pressure by inhibiting ACE activation [23]. Red ginger contains flavonoids, saponins, and non-flavonoid phenols. Flavonoids have an inhibitory effect on angiotensin-converting enzyme (ACE) activity [16] which reduces the formation of angiotensin II from angiotensin I, which results in vasodilation, reduces cardiac output and finally reduces blood pressure [17]. Inhibition of ACE can also increase nitric oxide and decrease superoxide anions, causing vasodilation [14]. Red ginger also contains phenolic compounds such as (6)-shogaol and (6)-gingerol, (10)-gingerol which have antioxidant effects [13]. Antioxidants are able to reduce free radicals such as thromboxane A₂, endothelins, and endoperoxides which are endothelial vasoconstriction factors [14, 16]. In addition, red ginger also contains saponins that play role inhibiting renin (RAA system) in the kidneys [16, 17]. so that reduces formation of angiotensin II and reduces aldosterone secretion which causes increase in excretion of salt and water by the kidneys and causes decrease in cardiac output [17].

When there is too much salt and water in the body, blood pressure increases by means of a complex physiological mechanism that alters venous return to the heart, causing an increase in cardiac output. Pathological conditions that change renal threshold will increase on systemic arterial pressure. Renin and angiotensin play a role in regulating blood pressure. Kidneys produce renin, an enzyme on the plasma protein substrate to secrete angiotensin I, which is converted by angiotensin converting enzyme into angiotensin II and then angiotensin III. Angiotensin II and III are potent vasoconstrictor of blood vessels and control the release mechanism of primary aldosteronism. In increased sympathetic nervous system activity, angiotensin II and III inhibitory effects on sodium excretion [16, 17]. Researchers argue that the compounds in red ginger lead to mechanism for lowering blood pressure in the vascular system (peripheral resistance) as vasodilator. Decreased peripheral resistance due to dilation of blood vessels, resulting in a decrease in systolic pressure without a decrease in diastolic pressure because there is no decrease in fluid (cardiac output) which is related to the mechanism of cardiac performance at diastolic pressure.

In S2 respondents, the control group, all respondents observed a decrease in systolic pressure, and 25 respondents observed a decrease in diastolic blood pressure (systolic $p = 0.72 > 0.05$ & diastolic $p = 0.600 > 0.05$) which means that giving placebo did not provide a significant change in blood pressure (Table 2). While the treatment group of all respondents observed a decrease in systolic pressure and 22 respondents observed a decrease in diastolic blood pressure (systolic $p = 0.000 < 0.05$ & diastolic $p = 0.000 < 0.05$), which means that the treatment of steeping binahong leaves gives a significant change in blood pressure (Table 3).

The flavonoids in binahong leaves are flavonols. These compounds can reduce oxidative stress, inhibit ACE activity, and increase relaxation of blood vessel endothelium [22]. Flavonoids inhibit the conversion of angiotensin I to angiotensin II, causing vasodilatation. Flavonoids as ACE inhibitors can also inhibit aldosterone secretion from the adrenal cortex, causes NaCl excretion will increase and finally blood pressure will decrease [21]. The difference in systolic and diastolic pressure results is due to the flavonoid content in binahong leaves which acts as an ACE inhibitor which can also improve arterial compliance and distensibility. This role can cause a decrease in systolic pressure that is more significant than diastolic [23]. Researchers argue that the

flavonoid mechanism, in this case Flavonol, in addition to providing a vasodilating effect, must also affect NaCl excretion through the RAAS system in the kidneys by inhibiting aldosterone secretion from the adrenal cortex, thereby reducing cardiac pre load, in this case it can be seen from the value diastolic. However, the results of this study did not show a significant decrease in the diastolic value, this may be because the amount of fluid that the respondent drank during the measurement process between the pre test and post test could not be controlled. These are the limitations and weaknesses of this study. So, it is necessary to do further research related to the mechanism of reducing blood pressure from steeping binahong leaves by adding urine parameters to assess the amount of fluid in each treatment condition.

4. Conclusions

Treatment of steeping red ginger rhizomes and steeping binahong leaves had a significant effect on reducing systolic blood pressure ($p = 0.78 > 0.05$).

5. References

- Sheps, S. G. 2005. Mayo Clinic Hipertensi, Mengatasi Tekanan Darah Tinggi. 2005. PT Intisari Mediatama. Jakarta.
- Yonata, A., & Pratama, A. S. P. Hipertensi sebagai faktor pencetus Terjadinya Stroke. *Jurnal Majority*. 2016. 5(3): 17-21.
- Park, J.B., Kario, K., dan Wang, J.G. Systolic Hypertension: An Increasing Clinical Challenge in Asia. *Hypertension Research*. 2015. Volume 38 (4): 227–236.
- Pusat data dan informasi kementerian kesehatan RI. 2014. Infodatin Hipertensi. Kementerian Kesehatan RI.
- Duke J, et al. 2000. Handbook of medicinal Herbs. CRC Press: United State of America.
- Kikuzaki, H. and N. Nakatani. Antioxidant effects of some ginger constituents. *J. Food Science*. 1993. 58: 1.407–1.410.
- Rahmah, N, A. Skrining Aktivitas Antihipertensi Dari Ekstrak Etanol 70% Rimpang: Jahe Merah (*Zingiber officinale* Roscoe), Bangle (*Zingiber purpureun* Roscoe), Temu Kunci (*Boesenbergia Rotunda*), Dan Temu Putih (*koempfria Rotunda*) Pada Tikus Yang Di Indikasikan Adrenalin. 2008. Jakarta: Program Studi Farmasi Fakultas Kesehatan UIN Syarif Hidayatullah.
- Zakaria, F.R. dan T.M. Rajab. Pengaruh ekstrak jahe (*Zingiber officinale* Roscoe) terhadap produksi radikal bebas makrofag mencit sebagai indikator imunostimulan secara in vitro. 1999. Persatuan Ahli Pangan Indonesia (PATPI): Prosiding Seminar Nasional Teknologi Pangan: 707–716.
- Mohan M Blalaraman, R. Kasture, SB. Antihypertensi activy *Zingiber officinale* ann korean gingseng in exprimentaly induced hypertension in rats. *Oriental pharmacy and Exprimental Medicine*. 2007. 7(3): 261-273.
- Suekawa, M, Ishige, A, Yuasa, K, Sudo, K, Aburada, M, Hosoya, E. Pharmacological studies on ginger I. Pharmacological actions of pungent constitutents, (6)-gingerol and (6)-shogaol. *Journal of Pharmacobio-Dynamics*. 1984. 7: 836-848.
- Suekawa, M, Aburada, M, Hosoya, E. Phamacological studies on ginger II. Pressor action of (6)-shogaolin anesthetized rats, orhindquarters, tail and mesenteric vascular beds of of rats. *Journal of pharmacobio-dynamic*. 1986. 9: 842-852
- Masuda, Y., Kikuzaki, H., Hisamoto, M., & Nakatani, N. Antioxidant properties of gingerol related compounds from ginger. *BioFactors*. 2004. 21(1–4), 293–296.

-
- Ghayur, M. N., & Gilani, A. H. Ginger Lowers Blood Pressure Through Blockade of Voltage. Dependent Calcium Channels. 2005. 45(1): 74–80.
- Kojsova, S., Jendekova, L., Zlcha, J., Kunes, J., Andriantsitohaina, R., & Pechanova, O. The Effect of Different Antioxidant on Nitric Oxide Production in Hypertensive Rats. *Physiological Research*. 2006.
- Katzung, B.G., Masters, S.B. dan Trevor, A.J. 2014. *Farmakologi Dasar & Klinik*, Vol.2 Edisi 12. Editor Bahasa Indonesia Ricky Soeharsono et al., Penerbit Buku Kedokteran EGC. Jakarta.
- Dipiro, T.J., wells, G.B, Schwinghammer, T.L. dan Dipiro C.V. 2012. *Pharmacotherapy Handbook Eight Edition*. The McGraw-Hill Companies. New York.
- H.H. Gray, K.D. Dawkins, J.M. Morgan, I.A. Simpson, *Kardiologi: Lecture Notes Ed 4*. 2005. Jakarta: Penerbit Erlangga.
- Astuti, S. M., Sakinah, M., Andayani, R. dan Risch, A. Determination of Saponin Compound from *Anredera cordifolia* (Ten) Steenis Plant (Binahong) to potential Treatment for Several Diseases. *Journal of Agricultural Science*. 2011. University Malaysia Pahang. Malaysia.
- Rachmawati, S. Studi Makroskopi dan Skrining Fitokimia Daun *Anredera cordifolia* (Ten.) Steenis. 2007. Fakultas Farmasi UNAIR. Surabaya.
- Sari, L. Pemanfaatan obat tradisional dengan pertimbangan manfaat dan keamanannya. *Maj. Ilmu Kefarmasian*. 2006. 3, 1–7.
- Santi Widiasari. Mekanisme Inhibisi Angiotensin Converting Enzym Oleh Flavonoid Pada Hipertensi. *Collaborative Medical Journal*. 2018. Universitas Abdurrah. Riau.
- Anggi Utri Aidasari. Aktivitas Antihipertensi Ekstrak Etanol Daun Binahong (*Anredera cordifolia* (Ten.) V. Steenis) dan Ekstrak Etanol Daun Tempuyung (*Sonchus arvensis* L.) beserta Kombinasi Pada Tikus Wistar Jantan yang Diinduksi Angiotensin II. 2018. Sains dan Teknologi Farmasi ITB. Bandung.
- Ghayur, M. N., Gilani, A. H., Afridi, B. M., & Houghton, J. P. Cardiovascular Effects of Ginger Aqueous Extract and Its Phenolic Constituents are Mediated Through Multiple Pathways. *Vascular Pharmacology*. 2005.

Comparison of blood pressure reduction between treatments of steeping Red Ginger Rhizome and steeping Binahong Leaves

GRADEMARK REPORT

FINAL GRADE

/0

GENERAL COMMENTS

Instructor

PAGE 1

PAGE 2

PAGE 3

PAGE 4

PAGE 5

PAGE 6

PAGE 7

PAGE 8

PAGE 9
