

# Immunomodulator Activity Test of Ethanol Extract of Sappan Wood (*Caesalpinia Sappan* L.) in Mice (*Mus Musculus*) Infected by *Staphylococcus Aureus*

*by Laily Latifah ,*

---

**Submission date:** 11-Sep-2022 06:58PM (UTC-0700)

**Submission ID:** 1897451436

**File name:** UAM-Martina\_Kurnia\_Rohmah.docx (107.53K)

**Word count:** 4312

**Character count:** 25443

**Immunomodulator Activity Test of Ethanol Extract of Sappan Wood (Caesalpinia Sappan L.) in Mice (Mus Musculus) Infected by Staphylococcus Aureus**

**Prueba De Actividad Inmunomoduladora Del Extracto Etanólico De Madera De Sappan (Caesalpinia Sappan L.) En Ratones (Mus Musculus) Infectados Por Staphylococcus Aureus**

Martina Kurnia Rohmah<sup>1a\*</sup>, Farida Anwari<sup>2b</sup>, Arif Rahman Nurdianto<sup>3b</sup>, Elisa Dwi Febrianti<sup>a</sup>,  
Juvita Anggraini<sup>a</sup>

ORCID ID: 0000-0003-4883-9609<sup>1</sup>

ORCID ID: 0000-0002-3796-0932<sup>2</sup>

ORCID ID: 0000-0002-3178-7112<sup>3</sup>

<sup>a</sup>Pharmacy, Faculty of Health Sciences, Universitas Anwar Medika, Sidoarjo, Indonesia

<sup>b</sup>Medical Laboratory Technology, Faculty of Health Sciences, Universitas Anwar Medika, Sidoarjo, Indonesia

\*Corresponding Author: Martina Kurnia Rohmah

Email: [martina.kurniarohmah@gmail.com](mailto:martina.kurniarohmah@gmail.com)

## SUMMARY

**Introduction:** The potential of immunomodulators from a natural product is widely studied as a choice of natural immune support supplements. In Indonesia, Caesalpinia sappan (Sappan wood) is a traditional herbal drink to boost the immune system. Sappan wood contains potential compounds as immunomodulators, such as brazilin, sappanchalcone, and phenol. This study aims to determine the immunomodulatory activity of Sappan wood infected by Staphylococcus aureus.

**Methods:** This study was a true experimental design that used Mus musculus infected by Staphylococcus aureus ( $10^{-9}$  CFU/ml). This study consisted of normal control, negative control (placebo), positive control (immune booster), and 96% ethanol extract of Sappan wood (SWEE) 25, 50, 100, and 200 mg/kg body weight. The treatment was given for 7 days after the Staphylococcus aureus infection.

**Results:** One-way ANOVA analysis showed that the administration of 96% ethanol extract from Sappan wood on phagocytic activity, C-Reactive Protein (CRP), and total leukocyte levels was significantly different from negative control ( $p=0,000$ ), but not to a normal and positive control ( $p<0,05$ ). The ethanol extract of Sappan wood at 100 and 200 mg/ kg body weight increases phagocytic index until 1,32 and 1,54 folds. It also can reduce CRP levels to 60 and 48 mg/L. The lower CRP level indicates lower inflammation.

**Conclusion:** Sappan wood has immunomodulatory activity through increased phagocytic activity, decreased CRP levels, and normalized total leukocyte levels, especially at 100 and 200 mg/kg body weight. The extract can enhance total leukocyte levels and positively affect CRP levels.

**Keywords:** C-Reactive Protein, Immunomodulator, Leukocytes, Phagocytosis, Sappan Wood

## RESUMEN

**Introducción:** El potencial de los inmunomoduladores de un producto natural es ampliamente estudiado como una opción de suplementos naturales de apoyo inmunológico. En Indonesia, Caesalpinia sappan (madera de Sappan) es una bebida tradicional a base de hierbas para estimular el sistema inmunológico. La madera de sappan contiene compuestos potenciales como inmunomoduladores, como brasilina, sappanchalcona y fenol. Este estudio tiene como objetivo determinar la actividad inmunomoduladora de la madera de Sappan infectada por Staphylococcus aureus.

**Métodos:** Este estudio fue un verdadero diseño experimental que utilizó Mus musculus infectado por Staphylococcus aureus ( $10^9$  UFC/ml). Este estudio consistió en un control normal, un control negativo (placebo), un control positivo (refuerzo inmunitario) y 25, 50, 100 y 200 mg/kg de peso corporal de extracto etanólico de madera de Sappan (SWEE) al 96 %. El tratamiento se administró durante 7 días después de la infección por Staphylococcus aureus.

**Resultados:** El análisis ANOVA de una vía mostró que la administración de extracto etanólico al 96 % de madera de Sappan sobre la actividad fagocítica, la proteína C reactiva (PCR) y los niveles totales de leucocitos fue significativamente diferente del control negativo ( $p=0,000$ ) pero no un control normal y positivo ( $p<0,05$ ). El extracto etanólico de madera de Sappan a 100 y 200 mg/kg de peso corporal aumenta el índice fagocítico hasta 1,32 y 1,54

veces. También puede reducir los niveles de CRP a 60 y 48 mg/L. El nivel más bajo de PCR indica una inflamación más baja.

**Conclusión:** La madera de sappan tiene actividad inmunomoduladora a través de una mayor actividad fagocítica, disminución de los niveles de PCR y normalización de los niveles totales de leucocitos, especialmente a 100 y 200 mg/kg de peso corporal. El extracto puede mejorar los niveles totales de leucocitos y afectar positivamente los niveles de PCR.

**Palabras Clave:** Proteína C Reactiva, Inmunomodulador, Leucocitos, Fagocitosis, Sappan Wood

## INTRODUCTION

The immune system has the basic function of protecting against foreign pathogens and infectious agents, consisting of innate and adaptive immunity with various cells and molecules involved (1–3). Innate immunity is a non-specific immune response as a first-line defence such as physical barriers, anatomical barriers, epithelial and phagocytic cell enzymes, phagocyte, inflammations-relate serum proteins, surface and phagocyte granule antimicrobial peptides, the immune receptor on cells, and cells that release cytokines and inflammatory mediators (4–6). Innate immunity is followed by adaptive immunity, a specific immune response and complex. Adaptive immunity consists of antibody response (B cell) and cell-mediated response (T helper and Cytotoxic T lymphocyte) (7).

Immunomodulators are synthetic or natural substances that can stimulate, suppress, or modulate any aspect of the immune system, including innate and adaptive immunity. In clinical medicine, immunomodulators are usually used to treat infection, reconstitute immunodeficiency, and suppress excessive immune function (8–10). Immunomodulators consist of immunoadjuvants, immunostimulants, and immunosuppressants. Immunoadjuvant is used for enhancing the efficacy of vaccines (specific immunostimulants). Immunostimulant is used to enhance the immune system and against infection. Immunosuppressant suppresses excessive immune function (11).

Currently, immunomodulator from natural compounds (phytochemical) is preferable because it has lower toxicity (12,13). The most common phytochemicals that can modulate immune response are flavonoids, flavanols, quinones, glycosides, polysaccharides, terpenoids, alkaloids, phenolics, saponins, various bitter, vitamin C, etc. (14).

Caesalpinia sappan (Sappan wood) is a traditional medicinal plant used in Asia, including Indonesia. Indonesians consume Sappan wood to increase stamina and maintain a healthy body. Sappan wood has antioxidant, antibacterial, antiviral, antifungal, cardiovascular protection, anti-inflammatory, antidiabetic, anticancer, anti-malarial, and gastroprotective roles (15). Sappan wood contains homoisoflavonoid and phenolics such as 4-O-methylsappanol, protosappanin A, protosappanin B, protosappanin E, brazilin, brazilein, caesalpin, brazilide A, neosappanone A, caesalpin P, sappanchalcone, 3-deoxysappanone, 10,7,3',4'-trihydroxy-3-benzyl-2H-chromene (16). Purified brazilin (10 mg/kg body weight) isolated from Sappan wood reduced the arthritis index score, and acute inflammation was administered every three days for 21 days. Other research shows that 124 mg Sappanchalcone from the dried heartwood of Sappan wood regulates the level of proinflammatory cytokines such as TNF- $\alpha$ , IL-1 $\beta$ , and IL-6 (17). In addition, Sappan wood (25 mg/kg body weight) has an immunomodulatory effect on murine peritoneal macrophages (18).

Staphylococcus aureus can cause some disturbances to cells that play a role in the immune system, one of which is the virulent Staphylococcus aureus in the form of A protein that can cause binding to immunoglobulin receptors (19). In addition, staphylococcus aureus causes excessive production of proinflammatory cytokines, resulting in inflammation and systemic shock (20). Leucocidin enzymes can destroy leukocytes, neutrophils and macrophages. Coagulase enzymes can accelerate the formation of fibrin so that it blocks the phagocytosis process. Staphylococcus aureus has also increased C-Reactive Protein levels (CRP) in vitro (21).

Leukocytes are an important part of the body's defence system, fighting infection-causing microorganisms, tumour cells, and harmful foreign substances. Leukocytes protect the body against various diseases by means of phagocytes and produce antibodies. The main cells that play a role in non-specific defence are mononuclear (monocytes and macrophages). Both types of cells are derived from white blood cells. Early phagocytosis of bacterial invasion can prevent disease (1).

C-Reactive Protein (CRP) is an alpha globulin present in the serum as a non-specific immune response when inflammation occurs. CRP is an acute phase protein that can increase in value up to 1000 times at the site of inflammation or infection. An immunomodulator is a drug that can be used to restore and repair an impaired immune system (22). Based on the above background, this research aims to determine the immunomodulatory activity of Sappan wood infected by Staphylococcus aureus.

## METHODS

This research was a true experimental design that used *Mus musculus* (mice) with pre and post-design. The mice were infected by *Staphylococcus aureus* ( $10^{-9}$  CFU/ml). This research was divided into 7 groups: normal control, negative control (CMC-Na 1%), positive control (immune booster), and 96% ethanol extract of Sappan wood 25, 50, 100, and 200 mg/kg body weight. The treatment was given for 7 days after the *Staphylococcus aureus* infection. This research is a pre-clinical study using experimental animals and has received an ethics certificate numbered 632/RSAM/III/2021.

The extraction of Sappan wood was carried out using the maceration method. The 500 grams of extracted sample was put into a vessel and dissolved in 3,750 ml of 96% ethanol as solvent. Maceration was carried out for 5 days with stirring 3 times a day. The extract was filtered to separate the filtrate. The filtrate was then concentrated with a rotary evaporator at a temperature of  $65^{\circ}\text{C}$  to obtain a thick extract of Sappan wood. Phytochemical tests were carried out to determine the content of Sappan wood extract, including flavonoids, alkaloids, tannins, polyphenols, anthraquinones, saponins, steroids, and terpenoids. Sappan wood ethanol extract solution was made with 4 concentrations: 25, 50, 100, and 200 mg/kg body weight. The treatment was adjusted according to the body weight of each mouse.

*Staphylococcus aureus* was grown on nutrient agar media and incubated for 1 x 24 hours. First, the preparation of  $10^{-9}$  CFU/ml bacterial suspension was done by adding 1 ml of bacterial suspension into 9 ml of sterile distilled water in the 1st test tube to form a bacterial concentration of  $10^{-1}$  CFU/ml. Then, the dilution was carried out by taking 1 ml in the 1st tube and adding 9 ml of sterilized distilled water to form a suspension with a concentration of  $10^{-9}$  CFU/ml.

Mice that had been acclimatized for 7 days were then infected with *Staphylococcus aureus*. After 1x24 hours, the mice were grouped into 7 test groups: normal control, negative control (CMC-Na 1%), positive control (immune booster), and 96% ethanol extract of Sappan wood 25, 50, 100, and 200 mg/kg body weight. The treatment was given for 7 days.

Parameters in this immunomodulatory activity test included total leukocyte count, phagocytosis index, and CRP levels. Measurement of leukocyte count, index, and CRP levels before and after treatment (pre and post-design). The total leukocyte was measured by using Hematology Analyzer. The phagocytosis index was measured using the Carbon Clearance method on a UV-Vis Spectrophotometer with a wavelength of 650 nm. CRP levels were

measured using a semi-quantitative method. The data that has been obtained is analyzed statistically using a t-test and one-way ANOVA.

## RESULTS

### Phytochemical Test of Sappan Wood Ethanol Extract

Based on the phytochemical test, it is known that the ethanol extract of Sappan wood contains the group of compounds shown in Table 1.

Table 1  
Phytochemical Test of Sappan Wood Ethanol Extract

Group of Compounds	Results
Flavonoid	+
Alkaloid	+
Polyphenol	+
Saponin	+
Anthraquinone	+
Steroid	-
Terpenoid	-

### Phagocytic Activity Test

Phagocytic activity was measured from the phagocytic index using the carbon clearance method. The result phagocytic activity test is shown in Table 2 and Figure 1.

Table 2  
Phagocytic Activity Test Pre and Post Design

Treatment Groups	Phagocytic Index	
	Pre (Mean ± SD)	Post (Mean ± SD)
Normal Control	1,4 ± 0,017	1,9 ± 0,008
CMC Na 1%/ infected only (Negative Control)	1,4 ± 0,003	1,6 ± 0,013
Immuno Booster (Positive Control)	1,6 ± 0,021	3,8 ± 0,015
SWEE 25 mg/ kg body weight	1,5 ± 0,011	2,1 ± 0,015



SWEE 50 mg/ kg body weight	1,7 ± 0,001	2,6 ± 0,003
SWEE 100 mg/ kg body weight	1,6 ± 0,008	2,9 ± 0,011
SWEE 200 mg/ kg body weight	1,7 ± 0,013	3,2 ± 0,007

SWEE: Sappan Wood Ethanol Extract

Based on paired sample t-test, post and pre-test is significantly different with significant value  $p=0,007$  ( $p<0,05$ ) and  $t\text{-count} > t\text{-table}$  ( $3,959 > 2,364$ ). This result shows that administration of Sappan wood ethanol extract increases phagocytic index.

The enhancement of the phagocytic index before and after treatment can be seen in the graph in Figure 1.




Figure 1

The Enhancement of Phagocytic Index Before and After Treatment. Treatment I: Normal Control, II: CMC Na 1%/ infected only (Negative Control), III: Immuno Booster (Positive Control), IV: SWEE 25 mg/ kg body weight, V: SWEE 50 mg/ kg body weight, VI: SWEE 100 mg/ kg body weight, VII: SWEE 200 mg/ kg body weight.

The graph shows that an immune booster can increase the phagocytic index 2.4 folds, while the administration of ethanol extract of Sappan wood 25, 50, 100, and 200 mg/ kg body weight successively increased the phagocytic index by 1.4, 1.5, 1.8 and 1.9 folds respectively. This result differs from the negative control group with *Staphylococcus aureus* infection only without treatment. The phagocytic index is quite low.



One-way ANOVA show there is significantly differences among 7 treatments with significant value  $p=0.000$  and  $F\text{-count} > F\text{ table}$  ( $639,48 > 2,29$ ). The result of one-way ANOVA is shown in Table 3.

Table 3  
The Difference in Phagocytosis Index Before and After Treatment

Treatment Groups	Mean $\pm$ SD
Normal Control	0,54 <sup>a</sup> $\pm$ 0,05
CMC Na 1%/ infected only (Negative Control)	0,18 <sup>b</sup> $\pm$ 0,04
Immuno Booster (Positive Control)	2,16 <sup>c</sup> $\pm$ 0,05
SWEE 25 mg/ kg body weight	0,6 <sup>a</sup> $\pm$ 0,1
SWEE 50 mg/ kg body weight	0,74 <sup>a</sup> $\pm$ 0,05
SWEE 100 mg/ kg body weight	1,32 <sup>d</sup> $\pm$ 0,04
SWEE 200 mg/ kg body weight	1,54 <sup>e</sup> $\pm$ 0,01

SWEE: Sappan Wood Ethanol Extract

The statistical test shows that the phagocytic index significantly differs between the infection condition (negative control) and the normal condition. The phagocytic index in normal conditions is higher than in infection conditions. The administration of an immune booster (positive control) is proven to increase phagocytic index 2,4 folds and is significantly different from the normal and negative control. The administration of ethanol extract of Sappan wood is proven to increase phagocytic index on 100 and 200 mg/ kg body weight until 1,32 and 1,54 folds. It is significantly different from normal and negative control.

### C-Reactive Protein (CRP) Test

The graph of CRP levels after each treatment is shown in Figure 2.




Figure 2

The CRP Level After Treatment and Staphylococcus aureus infection. Treatment I: Normal Control, II: CMC Na 1% infected only (Negative Control), III: Immuno Booster (Positive Control), IV: SWEE 25 mg/ kg body weight, V: SWEE 25 mg/ kg body weight, VI: SWEE 25 mg/ kg body weight, VII: SWEE 25 mg/ kg body weight.

CRP levels in the normal group, negative control, positive control, SWEE 25, 50, 100, and 200 mg/ kg body weight were 0, 96, 12, 96, 84, 60 and 48 mg/ L. CRP is not produced in the normal condition, but high in infection (negative control). The immune booster can decrease CRP level to 12 mg/ L. The ethanol extract of Sappan wood at 100 and 200 mg/ kg body weight can decrease CRP levels to 60 and 48 mg/L. The lower CRP level indicates lower inflammation.

### Total Leukocyte Level

The total leukocyte was measured on the last day after treatment. The total leukocyte is shown in Figure 3. The result shows that the normal condition's leukocyte level is  $\pm 10,5 \times 10^3$  cell/ mm<sup>3</sup>. Staphylococcus aureus infection increases leukocyte infection until  $\pm 16,7 \times 10^3$  cell/ mm<sup>3</sup>. The leukocyte level if positive control is  $\pm 11,2 \times 10^3$  cell/ mm<sup>3</sup>. The leukocyte ethanol extract of Sappan wood 25, 50, 100, and 200 mg/ kg body weight respectively are  $\pm 15,8 \times 10^3$ ,  $14,2 \times 10^3$ ,  $13,6 \times 10^3$ , and  $12,2 \times 10^3$  cell/ mm<sup>3</sup>. The leukocyte level of ethanol extract of Sappan wood is in the normal range.




Figure 3

The Leukocyte Level After Treatment and *Staphylococcus aureus* infection. Treatment I: Normal Control, II: CMC Na 1%/ infected only (Negative Control), III: Immuno Booster (Positive Control), IV: SWEE 25 mg/ kg body weight, V: SWEE 25 mg/ kg body weight, VI: SWEE 25 mg/ kg body weight, VII: SWEE 25 mg/ kg body weight.

## DISCUSSION

Based on the phytochemical test in this research, we know that Sappan wood contains flavonoid, alkaloid, polyphenol, saponin, and anthraquinone but not steroid and terpenoid. Based on the literature, ethanol 95% Sappan wood extract contains an alkaloid, flavonoid, anthraquinones, coumarin, saponin, tannin, and cardiac glycoside but does not contain steroid and terpenoid (23). Sappan wood also contains phenolic compounds, mainly including phenolic acids, flavonoids, tannins, coumarins, lignans, quinones, stilbenes, and curcuminoids (24). Sappan wood also contains several aromatic compounds such as brazilin, sappanchalcone, caesalpin, protosappanin A and B, homo-isoflavonoids  $\beta$ -sitosterol, mono-hydroxybrazilin, and quercetin (25).

Phagocytic tests show that ethanol extract of Sappan wood 100 and 200 mg/kg body weight significantly increases phagocytic index until 1,32 and 1,54 folds. Meanwhile, the immune booster (positive control) can increase the phagocytic index by 2,16 folds. Phagocytosis is a complex process for the ingestion and elimination of pathogens and apoptotic cells, tissue homeostasis, control the important aspect of inflammation and immune response. The phagocytic cells in the immune system such as monocytes, macrophages,

neutrophils, dendritic cells, osteoclasts, and eosinophils. Phagocytosis is a fundamental process in immunity (26). The CRP test showed that Sappan wood ethanol extract reduced inflammation with a lower level than the infection condition (negative control). Along with an increase in the phagocytic index and a decrease in inflammation after ethanol extract of Sappan wood treatment, the total leukocyte level gradually returns to normal after exceeding normal.

In this research, *Staphylococcus aureus* infection (negative control) shows that phagocytic index is lower than immune booster and treatment of Sappan wood. It is caused by the pathogenicity of *Staphylococcus aureus* can decrease phagocytosis. The virulence factor of this bacteria causes this pathogen to attach to a tissue's cells, escape the host immune system, and have some factors that decrease phagocytosis, factors that interact with anti-staphylococcal antibodies, and factors that elaborate proteases, exotoxins, and enzyme in the immune system. *Staphylococcus aureus* is a pyogenic pathogen capable of tissue invasion and evasion of phagocytosis by neutrophils. Staphylococcal lipoteichoic acid is involved in the synthesis of inflammatory cytokines by monocyte/ macrophages (27). *Staphylococcus aureus* can activate pro-inflammatory cytokines, including IL-6 and IL-8 (28).

The previous research showed that 25 mg/kg body weight of ethanol extract of *Caesalpinia sappan* had immunomodulatory activity on peritoneal macrophage of albino mice. It significantly enhanced the phagocytic activity of macrophages (18). *Caesalpinia sappan* extract dose-dependently inhibited the expression of proinflammatory cytokines IL-1 $\beta$  and TNF- $\alpha$  in IL-1 $\beta$ -stimulated chondrocytes and LPS-stimulated THP-1 macrophages (29). Some of the compounds in Sappan wood can act as immunomodulators, such as polyphenol, flavonoid, quercetin, brazilin, coumarin, episappanol, protosappanin C, brazilin, isoprotosappanin B, and sappanol.

This research shows that ethanol extract of Sappan wood contains polyphenol and flavonoids. Polyphenol successfully isolated from Sappan wood is sappanchalcone, caesalpinia phenol G, and quercetin (30). Polyphenol can increase phagocytosis and optimize macrophage function (31).

The previous research shows that levels of flavonoids in Sappan wood extract using the maceration method is 0,0539% (32). The total flavonoid content correlates with the phagocytic capacity of macrophages (33). Flavonoids are also anti-inflammatory in several chronic diseases such as cancer, diabetes mellitus, cardiovascular disease, and neuroinflammation (34).

Quercetin is a type of flavonoid that has immunomodulatory activity. Quercetin promotes some gene expressions involved in phagocytosis, such as CORO1A, CYBA, LAMP1, RAB7A, RAC1 and PAK1 genes (35). In addition, quercetin inhibits the production of inflammation-producing enzymes such as cyclooxygenase (COX) and lipoxygenase (LOX), inhibit proinflammatory cytokine production against H<sub>2</sub>O<sub>2</sub>-induced inflammation, and suppresses many inflammatory pathways (36).

Brazilin, the major compound of Sappan wood, was reported to have anti-inflammatory. Brazilin suppressed the release of IL-1 $\beta$ , TNF- $\alpha$ , NO, and PGE<sub>2</sub>, suggesting that these effects are mediated by Heme oxygenase-1. This result correlates well with studies on Caesalpinia sappan extract and brazilin in mouse macrophages, demonstrating that the potential anti-inflammatory effects of the agents involve the inhibition of PGE<sub>2</sub> production (37).

Several compounds that have other immunomodulatory effects in Sappan wood include coumarin, episappanol, protosappanin C, brazilin, isoprotosappanin B, and sappanol. Coumarin increases phagocytosis activity. Episappanol, protosappanin C, brazilin, isoprotosappanin B, and sappanol significantly inhibited the secretion of the proinflammatory cytokines such as interleukin (IL-6) and tumor necrosis factor-alpha (TNF- $\alpha$ ). Sappanol increased the secretion of the anti-inflammatory IL-10 (29).

## CONCLUSION

In conclusion, Sappan wood ethanol extract significantly increases phagocytic index and suppresses C-reactive protein (inflammatory acute-phase reactant). The phagocytic index and CRP level of 100 and 200 mg/kg body weight ethanol extract of Sappan wood were significantly different from negative control (infection). This condition indicates the work of the immune system and inflammation was quickly completed so that the total leukocytes on the 7<sup>th</sup> day also gradually return to normal.

## REFERENCES

1. Nicholson LB. The immune system. *Essays Biochem.* 2016;60(3):275–301.
2. Wahyuningsih SPA, Pramudya M, Putri IP, Winarni D, Savira NII, Darmanto W. Crude polysaccharides from okra pods (*abelmoschus esculentus*) grown in Indonesia enhance the immune response due to bacterial infection. *Adv Pharmacol Sci* [Internet].

2018;2018. Available from: <https://www.scopus.com/inward/record.uri?eid=2-s2.0-85056608618&doi=10.1155%2F2018%2F8505383&partnerID=40&md5=ec3a241bd81d228281664ec6706dd563>

3. Jariah ROA, Hakim MS. Interaction of phages, bacteria, and the human immune system: Evolutionary changes in phage therapy. *Rev Med Virol*. 2019;29(5):e2055.
4. Levy ARRA, Rojas-villarraga A, Levy RA. Cancer and Autoimmunity. *Cancer and Autoimmunity*. 2000.
5. Indrawati R, Luthfi M, Oki A, Yuliati Y, Sosiawan A, Rachmadi P, et al. Analysis of interleukin-10 anti-inflammatory cytokines in salivary lymphocyte surface: A cross sectional study. *J Int Oral Heal* [Internet]. 2020;12(5):439–42. Available from: [https://www.scopus.com/inward/record.uri?eid=2-s2.0-85095720915&doi=10.4103%2Fjih.jih\\_293\\_18&partnerID=40&md5=76201dd6777fb7280ba5ebd1ccaf5fe3](https://www.scopus.com/inward/record.uri?eid=2-s2.0-85095720915&doi=10.4103%2Fjih.jih_293_18&partnerID=40&md5=76201dd6777fb7280ba5ebd1ccaf5fe3)
6. Etika R, Sudarmo SM, Suwarno S, Mapindra MP, Mahindra MP. Analytical study of inflammatory cytokines and immunoglobulin expression following dust mite allergen exposure in pregnant mice. *Paediatr Indones*. 2021;61(6):336–42.
7. CHAFFEY N. Alberts, B., Johnson, A., Lewis, J., Raff, M., Roberts, K. and Walter, P. *Molecular biology of the cell*. 4th edn. Ann Bot. 2003;91(3):401–401.
8. Shantilal S, S. VJ, B SSS. Review On Immunomodulation And Immunomodulatory Activity Of Some Medical Plant. *Eur J Biomed Pharm Sci*. 2018;5(8):163–74.
9. Sabdoningrum EK, Hidanah S, Ansori ANM, Fadholly A. Immunomodulator y a nd Antioxidant Activities o f Phyllanthus niruri L. Extract a gainst t he Laying Hens Infected b y Escherichia c oli. *Res J Pharm Technol*. 2020;13(5):2246–50.
10. Pramudya M, Wahyuningsih SPA. Immunomodulatory potential of polysaccharides from coriolus versicolor against intracellular bacteria neisseria gonorrhoeae. *Vet World* [Internet]. 2019;12(6):735–9. Available from: <https://www.scopus.com/inward/record.uri?eid=2-s2.0-85067282382&doi=10.14202%2Fvetworld.2019.735-739&partnerID=40&md5=48a83111efe279fae2afbbfda39de6f1>
11. Sharma R, Rohilla A, Arya V. A short review on pharmacology of plant immunomodulators. *Int J Pharm Sci Rev Res*. 2011;9(2):126–31.
12. Di Sotto A, Vitalone A, Di Giacomo S. Plant-derived nutraceuticals and immune system modulation: An evidence-based overview. *Vaccines*. 2020;8(3):1–34.
13. Setiyowati PAI, Solekha R, Negara SBSMK, Rosalina R. Immunomodulator Effect of

- Lemongrass Extract (*Cymbopogon nardus* L.) to Increase Immune Cells as a Precaution Against SARS-CoV-2. *Biomol Heal Sci J*. 2021;4(2):73–7.
14. Nagoba B, Davane M. *iMedPub Journals Natural Immunomodulators*. *J Immunol Microbiol*. 2018;2(March):3–4.
  15. K. VA. *Caesalpinia sappan* L.: REVIEW ARTICLE. *Proc Int Conf Appl Sci Heal*. 2019;(4):290–9.
  16. Namikoshi M, Nakata H, Saitoh T. Homoisoflavonoids and Related Compounds. V.1) A Novel Dibenzoquinone Derivative from *Caesalpinia sappan* L. *Chem Pharm Bull*. 1987;35(9):3615–9.
  17. Jung EG, Han K Il, Hwang SG, Kwon HJ, Patnaik BB, Kim YH, et al. Brazilin isolated from *caesalpinia sappan* L. inhibits rheumatoid arthritis activity in a type-II collagen induced arthritis mouse model. *BMC Complement Altern Med*. 2015;15(1):1–11.
  18. Sunitha VS, Sunil MA, Radhakrishnan EK, Mathew J. Immunomodulatory Activity of *Caesalpinia sappan* L. Extracts on Peritoneal Macrophage of Albino Mice. *Int J Sci Res*. 2015;4(12):449–52.
  19. Kobayashi SD, DeLeo FR. Staphylococcus aureus protein A promotes immune suppression. *MBio*. 2013;4(5):4–6.
  20. Xu SX, McCormick JK. Staphylococcal superantigens in colonization and disease. *Front Cell Infect Microbiol*. 2012;2(April):52.
  21. Mölkänen T, Ruotsalainen E, Rintala EM, Järvinen A. Predictive Value of C-Reactive Protein (CRP) in identifying fatal outcome and deep infections in staphylococcus aureus bacteremia. *PLoS One*. 2016;11(5):1–14.
  22. Sproston NR, Ashworth JJ. Role of C-reactive protein at sites of inflammation and infection. *Front Immunol*. 2018;9(APR):1–11.
  23. Hemthanon T, Ungcharoenwiwat P. Antibacterial activity, stability, and hemolytic activity of heartwood extract from *Caesalpinia sappan* for application on nonwoven fabric. *Electron J Biotechnol*. 2022;55:9–17.
  24. Cai Y, Luo Q, Sun M, Corke H. Antioxidant activity and phenolic compounds of 112 traditional Chinese medicinal plants associated with anticancer. *Life Sci*. 2004;74(17):2157–84.
  25. Mehta PP, Shah RM, Shinde VM, Kamble RN, Mahadik KR. Phytochemical and pharmacological aspects of sandalwood. *Indian Drugs*. 2014;51(10):5–15.
  26. Rosales C, Uribe-Querol E. Phagocytosis: A Fundamental Process in Immunity.



Biomed Res Int. 2017;2017.

27. Fournier B, Philpott DJ. Recognition of *Staphylococcus aureus* by the innate immune system. *Clin Microbiol Rev.* 2005;18(3):521–40.
28. Ngo QV, Faass L, Sähr A, Hildebrand D, Eigenbrod T, Heeg K, et al. Inflammatory Response Against *Staphylococcus aureus* via Intracellular Sensing of Nucleic Acids in Keratinocytes. *Front Immunol.* 2022;13(February):1–12.
29. Mueller M, Weinmann D, Toegel S, Holzer W, Unger FM, Viernstein H. Compounds from *Caesalpinia sappan* with anti-inflammatory properties in macrophages and chondrocytes. *Food Funct.* 2016;7(3):1671–9.
30. Nguyen VB, Vu BD, Pham GK, Le BQ, Nguyen VC, Men C Van, et al. Phenolic Compounds from *Caesalpinia sappan*. *Pharmacogn J.* 2020;12(2):410–4.
31. Da Cunha LR, Muniz-Junqueira MI, Dos Santos Borges TK. Impact of polyphenols in phagocyte functions. *J Inflamm Res.* 2019;12:205–17.
32. Dwi Rusita Y. Flavonoids content in extracts secang (*Caesalpinia Sappan L.*) maceration method infundation analysis and visible ultraviolet spectrophotometer. *Int J Med Res Heal Sci.* 2016;5:176–81.
33. Munawaroh R, Prawita Setyowati E, Murwanti R, Hertiani T, Mada G, Penelitian Lingkungan Hidup dan Kehutanan Kupang B. Correlation Between Total Flavonoid Contents and Macrophage Phagocytosis Activity of Fractions From Faloak (*Sterculia quadrifida R.Br.*) Barks Ethanolic Extract In Vitro Korelasi Antara Kadar Flavonoid Total dan Aktivitas Fagositosis Makrofag Fraksi-Fraksi. *Tradit Med J.* 2018;23(1):47–55.
34. Ginwala R, Bhavsar R, Chigbu DGI, Jain P, Khan ZK. Potential role of flavonoids in treating chronic inflammatory diseases with a special focus on the anti-inflammatory activity of apigenin. *Antioxidants.* 2019;8(2):1–28.
35. Srikok S, Nambut S, Wongsawan K, Chuammitri P. Quercetin promotes the expression of genes involved in phagocytosis in bovine neutrophils. *Am J Anim Vet Sci.* 2017;12(2):85–95.
36. Li Y, Yao J, Han C, Yang J, Chaudhry MT, Wang S, et al. Quercetin, inflammation and immunity. *Nutrients.* 2016;8(3):1–14.
37. Lin FR, Niparko JK, Ferrucci and L. 基因的改变 NIH Public Access. *Bone.* 2014;23(1):1–7.



# Immunomodulator Activity Test of Ethanol Extract of Sappan Wood (*Caesalpinia Sappan* L.) in Mice (*Mus Musculus*) Infected by *Staphylococcus Aureus*

## ORIGINALITY REPORT

16%

SIMILARITY INDEX

13%

INTERNET SOURCES

13%

PUBLICATIONS

3%

STUDENT PAPERS

## PRIMARY SOURCES

1	<a href="http://www.researchgate.net">www.researchgate.net</a> Internet Source	2%
2	Shengqian Q. Wu, Miguel Otero, Frank M. Unger, Mary B. Goldring et al. "Anti-inflammatory activity of an ethanolic <i>Caesalpinia sappan</i> extract in human chondrocytes and macrophages", <i>Journal of Ethnopharmacology</i> , 2011 Publication	2%
3	Submitted to Universitas Pendidikan Indonesia Student Paper	2%
4	<a href="http://cmr.asm.org">cmr.asm.org</a> Internet Source	1%
5	<a href="http://jifi.farmasi.univpancasila.ac.id">jifi.farmasi.univpancasila.ac.id</a> Internet Source	1%
6	<a href="http://link.springer.com">link.springer.com</a> Internet Source	1%

7	<a href="http://apjr.net">apjr.net</a> Internet Source	1 %
8	<a href="http://www.chemfaces.com">www.chemfaces.com</a> Internet Source	1 %
9	<a href="http://bmccomplementmedtherapies.biomedcentral.com">bmccomplementmedtherapies.biomedcentral.com</a> Internet Source	1 %
10	<a href="http://uknowledge.uky.edu">uknowledge.uky.edu</a> Internet Source	1 %
11	<a href="http://repo.unand.ac.id">repo.unand.ac.id</a> Internet Source	1 %
12	<a href="http://journal.fk.unpad.ac.id">journal.fk.unpad.ac.id</a> Internet Source	1 %
13	TA Del-Valls, JM Forja, A Gomez-Parra. "The use of multivariate analysis to link sediment contamination and toxicity data to establish sediment quality guidelines an example in the Gulf of Cadiz (Spain)", Ciencias Marinas, 2007 Publication	1 %
14	Yuexing Chang, Ailing Guo, Yanlin Jing, Junjie Lin, Yuanyuan Sun, Lulu Kong, Haohang Zheng, Yun Deng. "Immunomodulatory activity of puerarin in RAW264.7 macrophages and cyclophosphamide-induced immunosuppression mice", Immunopharmacology and Immunotoxicology, 2021	1 %

15

studyres.com

Internet Source

1 %

---

16

Submitted to Leiden University

Student Paper

1 %

---

17

microbiologyjournal.org

Internet Source

1 %

---

Exclude quotes      On

Exclude matches      < 1%

Exclude bibliography      On