

TRYGLSERIDE LEVEL ON COFFEE CONSUMPTION

by Dora Turista

Submission date: 29-Oct-2023 08:22PM (UTC+0700)

Submission ID: 2210519520

File name: Coffee_and_Trigliseride_R2_1.docx (109.42K)

Word count: 3707

Character count: 21420



JURNAL TEKNOLOGI LABORATORIUM




Journal Homepage: www.teknolabjournal.com
ISSN 2580-0191(Online) | ISSN 2338 – 5634(Print)

Original Research



TRYGLESERIDE LEVEL ON COFFEE CONSUMPTION



Amellya Octifani^{1*}, Tarisa Suci Novianti¹,
Farida Anwari¹, Arif Rahman Nurdianto^{1*}, Fery Setiawan^{2,3},
Rizal Fauzi Nurdianto^{4,5}

- ¹ D3 Teknologi Laboratorium Medis, Fakultas Ilmu Kesehatan, Universitas Anwar Medika
- ² Departemen Patologi Mulut dan Maksilofasial, Fakultas Kedokteran Gigi, Universitas Airlangga, Indonesia
- ³ Doctoral Program, Faculty of Medicine, Universitas Airlangga, Surabaya, Indonesia
- ⁴ Graduate Student, Fakultas Kedokteran, Universitas Wijaya Kusuma, Indonesia
- ⁵ Magister Hukum Kesehatan, Universitas Soegijapranata, Indonesia

Abstract: Coffee is a drink that is commonly consumed by Indonesian people. Coffee contains caffeine, cafestol, and kahweol which can increase lipid levels, including increasing triglyceride levels. Triglycerides are substances consisting of glycerol linked to fatty acid groups. The purpose of this study was to describe blood triglyceride level in coffee addict students at Anwar Medika University using the GPO-PAP method. This study uses a quantitative descriptive design with a cross-sectional survey approach. This research was conducted in May 2023. The research sample consisted of 48 samples which were determined using a purposive sampling technique which was taken at random. The univariate analysis used is the distribution of frequencies and percentages which describe the presentation of data for one variable. This research test obtained an average triglyceride level of 158.7 mg/dl. Based on age category, maximum triglyceride levels were 204.73 mg/dl with an average of 152.26 mg/dl. Based on lifestyle, the maximum triglyceride level is 189.93 mg/dl with an average of 166.43 mg/dl. Based on the frequency of coffee consumption, the maximum triglyceride level is 160.64 mg/dl with an average of 158.51 mg/dl. Based on physical activity, the maximum triglyceride level is 173.88 mg/dl with an average of 160.60 mg/dl. Based on the type of coffee consumed, the maximum triglyceride level was 166.76 mg/dl with an average of 164.57 mg/dl. So, it can be concluded that excessive coffee consumption can increase triglyceride levels in the body.

Keywords: Coffee; Triglycerides; College Students.

INTRODUCTION

Coffee is a drink commonly consumed by Indonesian people. The distinctive aroma of coffee has its attraction for consumption. Indonesia is the third largest coffee-producing country in the world after Brazil and Vietnam with a total production of 748 thousand tons or 6.6% of world coffee production in 2016. From 2016 to 2019 the level of coffee consumption in Indonesia increased every year. It is predicted that this increase will increase by an average of 8.22% every year¹. The increase in coffee consumption in Indonesia is due, in part, to people's lifestyles which have made processed coffee drinks a daily activity for their daily

needs. However, the amount of coffee consumption in Indonesia still reaches 300 thousand tons and is still far below other countries².

The habit of drinking coffee has spread and become a culture in various regions, including among students, from urban to rural areas on the island of Java. Students also have a large level of coffee consumption due to internal and external factors. One of them is students who work while studying with heavy physical activity, then need coffee to prevent drowsiness and refresh the body. This causes the level of coffee consumption among young people to remain very high³.

The ingredients contained in coffee consist of more than a thousand molecules of different substances, including phenolic compounds, vitamins, minerals, and alkaloids which can increase lipid levels. The caffeine, cafestol, and kahweol compounds contained in coffee can also increase triglyceride levels⁴. Kalel et al.'s 2020 research on the effect of coffee consumption on serum lipid profiles in adults concluded that individuals who drink coffee show a higher risk of developing dyslipidemia⁵.

Triglycerides are a type of fat that is transported in the blood and stored in the body's fat tissue. Triglycerides are the main constituent of lipids in the body. Triglycerides are used by the body to provide energy in metabolic processes, small amounts of triglycerides are also used throughout the body to form cell membranes. Triglyceride levels are an indicator of body health, excessive triglycerides in the body can narrow blood vessels and increase the risk of heart attack⁶. Several factors that can influence triglyceride levels include age, gender, lifestyle, frequency of coffee consumption, hormone levels, and obesity. Triglyceride levels in the body can be controlled with a healthy lifestyle and avoiding foods or drinks that can increase lipid levels⁷.

This research aims to determine the description of triglyceride levels in the blood of students who are coffee addicts at Anwar Medika University using the GPO-PAP method.

MATERIAL AND METHOD

This research uses a quantitative descriptive method with a cross-sectional survey approach. The sampling technique in this research was purposive sampling which was taken randomly. Triglyceride examination uses the GPO-PAP (Glycerol Peroxidase Phosphate Acid) enzymatic colorimetric method. The principle of the GPO-PAP method is that triglycerides will be enzymatically hydrolyzed as glycerol and free acids with specific lipase to form a colored complex whose levels can be measured using a photometer. The intensity of the color formed can be determined by measuring the absorbance in the wavelength range 480-550 nm^{8,9}. This research was carried out in May 2023, at the Medical Biology Laboratory on the 4th floor of Anwar Medika University.

The tools used in this research were a tourniquet, red vacuum tube, 3cc syringe, micro lab 300 photometer, yellow and blue tip, 500-1000 ul and 50-5 ul micropipette, serology tube, and serology tube rack. The materials used in this research were 70% alcohol, blood serum/plasma samples, GPO-PAP triglyceride reagent, tissue, and labels/labels.

The procedure of this research was:

1. Pre-analytical, including sample preparation, labeling, venous blood sampling,
2. Analytical, including making serum/plasma and checking triglyceride levels as follows:

	Blanko	Test
Reagen	1000ul	1000µl

Sample	-	10 μ l
--------	---	------------

- Pipette 1000 μ l of triglyceride reagent with a micropipette into a serology tube then pipette 10 μ l of serum sample.
 - Homogenize, then incubate for 5-10 minutes
 - Read on the Microlab 300 instrument and record the results.
3. Post-analytic, including interpreting the results according to the reference value for triglyceride examination and recording the results then continuing with data analysis

The data analysis used in this research is manually using a Microsoft Excel computer with univariate analysis to analyze each variable from a study. The univariate analysis used is frequency and percentage distribution which describes the presentation of data for one variable¹⁰.

RESULTS AND DISCUSSION

Table 1 Research Subject Characteristic

Characteristic	Total	
	N	%
Age (tahun)		
18-19	9	17
20-21	13	23
22-23	11	26
24-25	15	34
Drinking Coffee Frequency		
< 3 glasses/day	23	49
> 3 glasses/day	25	51
Physics Activities		
Usually, physics activities	30	47
Seldom physics activities	18	53
Lifestyle		
Smoker	16	57
Non-Smoker	32	43
The amount of triglycerides		
Normal <150 mg/dl	28	58
High >150 mg/dl	20	42

Based on Table 1, the characteristics of the 48 research subjects based on triglyceride levels were divided into the normal category (<150 mg/dl) for 48 (58%) respondents and the high category (\geq 150 mg/dl) for 28 (42%) respondents. Age characteristics are categorized as ages ranging from 18-25 years with a frequency of coffee consumption of <3 cups/day and >3 cups/day as well as physical activity and lifestyle which are categorized into groups of regular exercise, rarely exercise, smoking, and non-smoking.

Table 2 Statistic Distribution of the amount of Triglycerides

Variable	N	Mean	SD	Min	Max
The amount of triglycerides (mg/dl)	48	158.7	67.089	62	304

Based on Table 2, the results of the statistical analysis of triglyceride levels in this study showed that the average triglyceride level was 158.7 mg/dl with the lowest triglyceride level being 62 mg/dl and the highest triglyceride level being 304 mg/dl. This can be caused by factors such as age, physical activity, frequency of daily coffee consumption, lifestyle, and the type of coffee consumed. Increasing age affects the decline in the function of the hormones estrogen and testosterone in distributing fat, thus allowing fat to accumulate in the body⁷. Consuming coffee over a long period can also cause an increase in lipids in the body, the compounds in coffee can speed up the process of narrowing and blockage of blood vessels which function to carry oxygen to the heart. Excessive levels of caffeine, cafestol, and kahweol compounds in the body will disrupt and inhibit triglyceride metabolism in the blood, causing lipid build up in blood vessels. Free fatty acids released due to excessive lipid accumulation can inhibit lipogenesis and will inhibit serum triacylglycerol clearance, resulting in increased blood triglyceride levels or hypertriglyceridemia^{11,12}.

Examination of triglyceride levels is an important parameter in the lipid profile which can help diagnose a disease or determine risk factors for coronary heart disease and detect metabolic syndrome, as well as monitor the effectiveness of lipid-lowering therapy. Based on analysis of the results of research conducted by researchers, an overview of triglyceride levels in coffee addicted students at Anwar Medika University was obtained with an average triglyceride level of 158.7 mg/dl. This can be caused by factors such as age, physical activity, frequency of daily coffee consumption, lifestyle and the type of coffee consumed. Consuming coffee over a long period of time can cause an increase in lipids in the body, the compounds in coffee can speed up the process of narrowing and blockage of blood vessels which function to carry oxygen to the heart. The habit of consuming coffee is closely related to the emergence of lipid disorders, including increased triglyceride levels¹³.

Table 3 The Distribution of the amount of triglyceride based on the coffee type

Coffee Type	The amount of triglyceride		Mean of triglyceride (mg/dl)
	<150 mg/dl	≥150 mg/dl	
Black coffee	10	9	166.76
Sachet coffee	13	16	162.38
Total	48		164.57

Based on Table 3, the results of the analysis of triglyceride levels in students who are coffee addicts based on the type of coffee consumed, namely black coffee and sachet coffee, showed that the average triglyceride level was 164.57 mg/dl. black coffee and sachet coffee. both contain caffeine, cafestol, and kahweol, but in different amounts or levels.^{14,15} Consuming black coffee tends to have higher triglyceride levels compared to those consuming sachet coffee.^{16,17} This is because black coffee has a caffeine content of 85-185 mg/cup, kahweol 6-12 mg/cup, and cafestol 4-6 mg/cup. Sachet or instant coffee has a caffeine content of 30-90 mg/coffee sachet in 150 ml of water, kahweol 0.2-0.6 mg/cup, and cafestol 0.1 mg/cup^{4,7,18,19}.

Triglycerides in the body will be converted into fatty acids and glycerol which are stored in adipose tissue and then absorbed through the intestines and distributed widely throughout the body. Excessive free fatty acids in the blood,

some will be used as an energy source and some will be taken to the liver as raw material for the formation of triglycerides. Free fatty acids will become triglycerides again and become part of VLDL (Very Low-Density Lipoprotein) because there has been a process of inhibiting the beta oxidation mechanism in the liver so that the lipids produced will be very rich in triglycerides. Excessive levels of caffeine, cafestol and kahweol compounds in the body will disrupt and inhibit triglyceride metabolism in the blood, causing lipid buildup in blood vessels. Free fatty acids released due to excessive lipid accumulation can inhibit lipogenesis and will inhibit serum triacylglycerol clearance, resulting in an increase in blood triglyceride levels or hypertriglyceridemia. In addition, the accumulation of lipids will cause adipose cells to be unable to store triglycerides adequately, which will trigger an increase in LDL (Low Density Lipoprotein) and ultimately an increase in triglyceride levels in the body. Caffeine is a central nervous system stimulant which can increase heart rate and contribute to the occurrence of supraventricular tachycardia (heart rhythm disturbance). The caffeine compound content in coffee has the function of stimulating nervous system activity and increasing heart function, but if caffeine is consumed in excess, caffeine will be toxic by inhibiting nervous system mechanisms and can increase triglyceride levels. Kahweol is a compound usually found in coffee that can cause the degradation of toxic substances and is protective against aflatoxin B1 if consumed in excess. The cafestol compound contained in coffee can also increase triglyceride levels by inhibiting the beta oxidation mechanism, preventing the breakdown of triglycerides into energy so that triglyceride levels in the blood will increase²⁰⁻²².

The picture of triglyceride levels increasing in the 24-25 year age category is triglyceride levels. This is because with increasing age the function of the body's organs will decrease due to aging which is a risk factor for functional disorders. The age factor will make a person less physically active and increase the risk of developing more functional disorders compared to a younger age. Increasing age significantly increases the risk of degenerative diseases in both men and women. As we age, there will be a decline in the function of various body organs, making it difficult to achieve balance in triglyceride levels. An increase in lipids in the blood is associated with a decrease in the elimination of lipids as bile salts and a decrease in receptors that mediate the clearance process of LDL. This can result in triglyceride levels tending to increase more easily. Age factors influence the deterioration of body functions, including stiffness of blood vessels (shrinking and aging). In men, serum triglyceride concentrations increase to a peak in middle age, while in women they continue to rise until the age of 70 years. Increasing age also affects the decline in the function of the hormones estrogen and testosterone in distributing fat, thus allowing fat to accumulate in the body. The danger is that if this fat buildup sticks to the walls of blood vessels, it will narrow blood flow, especially if the blood vessels are old. This condition will result in blocked blood vessels and increased triglyceride levels which can cause a decrease in HDL^{23,24}.

The limitation of this research is that this research was carried out semi-quantitatively using a Microlab 300 type photometer so that the measurement results were less accurate because they were influenced by several factors, including the pipetting angle being less precise so that the ratio of reagent and blood volumes was not appropriate because pipetting was still done manually outside of the tool for reacting the sample with reagents so that it will affect the results of measuring falsely low or high triglyceride levels. The cleanliness factor of the tube can also influence the results of measuring triglyceride levels because the researchers did not use disposable tubes so it is possible that when washing the tube it was not clean enough so that soap residue was still attached to the walls of the tube which will affect the results of measuring triglyceride levels because triglyceride examination tends to be sensitive compared to lipid profile

examination. other. Contamination during the reaction or incubation process can also be a factor in the high or false results of measuring triglyceride levels because this research still uses semi-quantitative methods where the process is still carried out outside the equipment which allows contamination to occur. Future researchers are expected to be able to develop this research by using more sophisticated tools to obtain more accurate research results and avoid several limitations²⁵.

Control efforts that can be made to monitor triglyceride levels are by paying more attention to health, for example having regular check-ups at least once every 3 months to determine body condition including triglyceride levels, in addition to maintaining diet, lifestyle, and increasing physical activity and reducing the habit of consuming coffee every day. so that health problems do not occur, for example increasing triglyceride levels in the body^{12,13,26-28}.

CONCLUSION

Based on the research results obtained, it can be concluded as follows, examination of high triglyceride levels ≥ 150 mg/dl with a percentage of 58% and triglyceride levels which are classified as normal, namely < 150 mg/dl with a percentage of 42%. The average triglyceride level was 158.7 mg/dl with the lowest triglyceride level being 62 mg/dl and the highest triglyceride level being 304 mg/dl. High triglyceride levels in many respondents were caused by several factors including age, physical activity, frequency of daily coffee consumption, lifestyle, and type of coffee consumed. Excessive coffee consumption can increase triglyceride levels in the body.

AUTHORS' CONTRIBUTIONS

Amellya Octifani: designed the protocols, and executed the protocols. Tarisa Suci Novianti: prepared the samples, and data collection. Farida Anwari: data analytics and visualization statistically. Arif Rahman Nurdianto: wrote the draft manuscript. Fery Setiawan, Arif Rahman Nurdianto, and Rizal Fauzi Nurdianto: reviewed and supervised the manuscript, and wrote the final manuscript. All authors have read and approved the final manuscript.

ACKNOWLEDGEMENT

The author would like to thank Anwar Medika University for facilitating this research.

FUNDING INFORMATION

This research received no specific grant from any funding agency in the public, commercial, or not-for-profit sectors.

DATA AVAILABILITY STATEMENT

The utilized data to contribute to this investigation are available from the corresponding author upon reasonable request.

DISCLOSURE STATEMENT

The views and opinions expressed in this article are those of the authors and do not necessarily reflect the official policy or position of any affiliated agency of the authors. The data is the result of the author's research and has never been published in other journals.

REFERENCE

1. Luthfiah F, Gizi Poltekkes Kemenkes Mataram J, Prabu Rangkasari Dasan Cermen Mataram J, Wiwin Diarti M, Pauzi I, Rif S, Sabariah, ah. Kadar Kolesterol Total Pada Peminum Kopi Tradisional Di Dusun Sembung Daye

- Kecamatan Narmada Kabupaten Lombok Barat Total Cholesterol Levels In Traditional Coffee Drinkers Dusun Sembung Daye Narmada Lombok District West. *Jurnal Kesehatan Prima*. 2016; 10(1): 1101-1112.
2. Fajriana NH, Fajriati I, Kimia J, Sains F, Teknologi D, Islam U, Sunan N, Yogyakarta K. Analisis Kadar Kafein Kopi Arabika (*Coffea arabica* L.) Pada Variasi Temperatur Sangrai Secara Spektrofotometri Ultra Violet. *Analytical and Environmental Chemistry*. 2018; 3(2):148-162; <http://dx.doi.org/10.23960/aec.v3.i2.2018.p148-162>
 3. Latifa NH, Rochdiani DD. (2019). Kajian Strategi Pemasaran Usaha Kedai Kopi Kadaka Cafeteria Di Kota Bandung. *Jurnal Agribisnis dan Sosial Ekonomi Pertanian UNPAD*. 2019; 4(1):642-651; <https://doi.org/10.24198/agricore.v4i1.22913>.
 4. Darmayani S, Rosanty A, Rahmayani D, Kesehatan JA, Kesehatan P, Kendari K. Gambaran Kadar Kolesterol Total Pada Pecandu Kopi Kecamatan Poasia Kota Kendari Overview the Total Cholesterol Levels Of Coffee Addicts in Poasia Subdistrict, City Of Kendari. *Jurnal Penelitian*. 2018; 10(1):33-41; <https://doi.org/10.36990/hjip.v10i1.122>
 5. Kalel A, Almutairi A, Alyahiwi A, Jalaluddin A, Aljohani M, Aloufi R, Almatrafi R. (2020). Effect of coffee consumption on serum lipid profile among adult population in Madinah, Saudi Arabia. *International Journal of Medicine in Developing Countries*. 2020: 1527–1532. <https://doi.org/10.24911/ijmdc.51-1592983718>
 6. Gemilang B, Miro S. Hubungan Kadar Trigliserida dan Kolesterol-HDL Terhadap Kadar Alanine Aminotransferase pada Pasien Non Alcoholic Fatty Liver Disease. *Jurnal Kesehatan Andalas*. 2016; 5(1):179-184. <https://doi.org/10.25077/jka.v5i1.465>
 7. Susilawati F, Hk N, Keperawatan J, Tanjungkarang P. Faktor Resiko Kejadian Stroke Di Rumah Sakit. *Jurnal Keperawatan*. 2018;14(1): 41-48.
 8. Hardisari R, Koiriyah B. Gambaran Kadar Trigliserida (Metode Gpo-Pap) Pada Sampel Serum dan Plasma EDTA. *Jurnal Sains dan Teknologi Laboratorium Medik*. 2016;5: 27-31 www.teknolabjournal.com
 9. Arikunto S. Prosedur penelitian Suatu Pendekatan Praktik. Jakarta: Rineka Cipta. 2014
 10. Notoatmodjo. Metodologi Penelitian Kesehatan, Jakarta: Rineka Cipta. 2018
 11. Adi N, Jangga N, Isma F. Perbedaan Kadar Kolesterol dan Trigliserida Serum Dari Darah Yang Dibekukan Sebelum Disentrifus dan Yang Langsung Disentrifus. *Jurnal Media Analisis Kesehatan*. 2019; 10(2):171-178
 12. Febiola W, Kesehatan John Paul Pekanbaru AI, Akademi Kesehatan John Paul Pekanbaru MI. Hubungan Indeks Massa Tubuh (IMT) Terhadap Kadar Trigliserida Pada Wanita Usia 40-60 Tahun. *Jurnal Sains dan Teknologi Laboratorium Medik*. 2017; 2(1):2-7
 13. Hainun Nisa, Artha DE, Risma. Pengaruh Rokok Terhadap Kadar Kolesterol 2 Jam Setelah Merokok. *Jurnal Kesehatan Masyarakat*. 2013; 8(1): 33–38.
 14. Surma S, Romanczyk M, Zembala MO, Filipiak KJ. Coffee and lipid profile: from theory to everyday practice. *Folia Cardiologica*. 2023;18(1):24-30
 15. Wahyani AD, Kartini A. Perbedaan Kadar Trigliserida Serum Tikus *Srague Dawley* Pada Pemberian Kopi Robusta Filter dan Tanpa Filter. *Journal of Nutrition College*. 2012; 1(1):253-257
 16. Du Y, Lv Y, Zha W, Hong X, Luo Q. Effect of coffee consumption on dyslipidemia: A meta-analysis of randomized controlled trials. *Nutrition, Metabolism, and Cardiovascular Diseases*. 2020;30(12):2159-2170; <https://doi.org/10.1016/j.numecd.2020.08.017>
 17. Condon C, Rai A, Tse C, Briggs JD, Sarpong NAK, Sayed TE, Stanton A. 34 High intake of coffee positively correlated with total and LDL cholesterol in healthy young adults. *BMJ Journals*. 2018; 104:A25-A26; <http://dx.doi.org/10.1136/heartjnl-2018-ICS.34>

18. Rico DC, Rodriguez NB, Marin JC. Impact of Coffee Consumption on Lipid Profile and Dyslipidemia Risk: Protocol for an Umbrella Review. *Universitas Medica*. 2022;63(1):1-6
19. Shi L, Brunius C, Johansson I, Bergdahl IA, Rolandsson O, Guelpen Bv, Winkvist A, Hanhineva K, Landberg. Plasma metabolite biomarkers of boiled and filtered coffee intake and their association with type 2 diabetes risk. *The Association for the Publication of the Journal of Internal Medicine*. 2019; 287:405-421;doi:10.1111/joim.13009
20. Ilmiawati C, Fitri F, Rofinda ZD, Reza M. Green coffee extract modifies body weight, serum lipids and TNF- α in high-fat diet-induced obese rats. *BMC Research Notes*. 2020;13:208; <https://doi.org/10.1186/s13104-020-05052-y>
21. Lim D, Chang J, Ahn J, Kim J. Conflicting Effects of Coffee Consumption on Cardiovascular Diseases: Does Coffee Consumption Aggravate Pre-existing Risk Factors?. *Processes*. 2020;8(4):438; <https://doi.org/10.3390/pr8040438>
22. Chang HC, Nfor ON, Ho CC, Chen PH, Kung YY, Hsu SY, Tantoh DM, Liaw YC, Hsieh CF, Liaw YP. Changes in High-Density Lipoprotein Cholesterol Levels in Relation to Coffee Consumption Among Taiwanese Adults. *Journal of Multidisciplinary Healthcare*. 2020;13:1427-1432; <https://doi.org/10.2147/JMDH.S276395>
23. Castillo MDd, Gomez BF, Saez NM, DeHond AI, Martirosyan DM, Mesa MD. Coffee Silverskin Extract for Aging and Chronic Diseases. *Functional Foods for Chronic Diseases*. 2016:386-409
24. McAllister MJ, Waldman HS, Renteria LI, Gonzalez AE, Butawan MB, Bloomer RJ. Acute coffee ingestion with and without medium-chain triglycerides decreases blood oxidative stress markers and increases ketone levels. *Canadian Journal of Pshycology and Farmacology*. 2020;98(4):458; <https://doi.org/10.1139/cjpp-2019-0458>
25. Gebeheyu GM, Feleke DG, Molla MD, Admasu TD. Effect of habitual consumption of Ethiopian Arabica coffee on the risk of cardiovascular diseases among non-diabetic healthy adults. *Heliyon*. 2020;6:1-8; <https://doi.org/10.1016/j.heliyon.2020.e04886>
26. Nurdianto AR, Anhar CA, Anwari F, Charisma AM, Farida EA. Edukasi Pola Hidup Sehat dalam Mengontrol Kadar Glukosa Darah Puasa bagi Anggota Prolanis Puskesmas Trosobo, Sidoarjo saat Pandemi COVID – 19. *Prosiding Seminar Nasional Abdimas Ma Chung*. 2021: 389-399
27. Nurdianto AR, Setiawan F, Anwari F, Tena HAB, Nurdianto RF. *Imunologi Forensik dan Imunohematologi*. 2021.
28. Anhar CA, Prasetyorini T, Kuswandari F, Nurdianto AR. *Kimia klinik (faal liver, jantung, elektrolit, endokrin, dan gas darah)*. 2022

TRYGLESERIDE LEVEL ON COFFEE CONSUMPTION

ORIGINALITY REPORT

13%

SIMILARITY INDEX

10%

INTERNET SOURCES

10%

PUBLICATIONS

6%

STUDENT PAPERS

PRIMARY SOURCES

1	www.teknolabjournal.com Internet Source	3%
2	teknolabjournal.com Internet Source	3%
3	www.ijstr.org Internet Source	2%
4	Aziz Eghbali, Kazem Ghaffari, Roya shaykh Baygloo, Aygin Eghbali, Ali Ghasemi. "Polyneuropathy Associated with Age of Starting the Transfusion and Serum Ferritin Level in Iranian Patients with Thalassemia Major and Intermedia", Hemoglobin, 2023 Publication	1%
5	media.neliti.com Internet Source	1%
6	Submitted to Nexford University Student Paper	1%
7	&NA;. "Information for Critical Care Nurses", Dimensions of Critical Care Nursing, 2009 Publication	1%

8

www.ncbi.nlm.nih.gov

Internet Source

1 %

9

"Abstracts of the Asian Congress of Nutrition 2019", *Annals of Nutrition and Metabolism*, 2019

Publication

1 %

10

e-journal.unair.ac.id

Internet Source

1 %

Exclude quotes On

Exclude matches < 1%

Exclude bibliography On